

5.f Il trattamento riabilitativo nella spondilite anchilosante. Evidenze scientifiche

Angela Del Rosso, Susanna Maddali Bongi

- Altan L., Bingöl U., Aslan M., Yurtkuran M. (2006), *The effect of balneotherapy on patients with ankylosing spondylitis*, in «Scand. J. Rheumatol.», 35, pp. 283-289.
- Altan L., Korkmaz N., Dizdar M., Yurtkuran M. (2012), *Effect of Pilates training on people with ankylosing spondylitis*, in «Rheumatol. Int.», 32, pp. 2093-2099.
- Analay Y., Ozcan E., Karan A., Diracoglu D., Aydin R. (2003), *The effectiveness of intensive group exercise on patients with ankylosing spondylitis*, in «Clin. Rehabil.», 17, pp. 631-636.
- Braun J., Davis J., Dougados M., Sieper J., Van der Linden S., Van der Heijde D. (2006), *First update of the international ASAS consensus statement for the use of anti-TNF agents in patients with ankylosing spondylitis*, in «Ann. Rheum. Dis.», 65, pp. 316-320.
- Braun J., Van den Berg R., Baraliakos X., Boehm H., Burgos-Vargas R., Collantes-Estevez E. et al. (2011), *2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis*, in «Ann. Rheum. Dis.», 70, pp. 896-904.
- Casimiro L., Barnsley L., Brosseau L., Milne S., Robinson V.A., Tugwell P. et al. (2005), *Acupuncture and electroacupuncture for the treatment of rheumatoid arthritis*, in «The Cochrane Datab. Syst. Rev.», 19(4), CD003788.
- Ciprian L., Lo Nigro A., Rizzo M., Gava A., Ramonda R., Punzi L. et al. (2013), *The effects of combined SpA therapy and rehabilitation on patients with ankylosing spondylitis being treated with TNF inhibitors*, in «Rheumatol. Int.», 33, pp. 241-245.
- Colina M., Ciancio G., Garavini R., Conti M., Trotta F., Govoni M. (2009), *Combination treatment with etanercept and an intensive SpA rehabilitation program in active ankylosing spondylitis*, in «Int. J. Immunopathol. Pharmacol.», 22, pp. 1125-1129.
- Dagfinrud H., Kvien T.K., Hagen K.B. (2008), *Physiotherapy interventions for ankylosing spondylitis*, in «The Cochrane Datab. Syst. Rev.», 23(1), CD002822.
- Dundar U., Solak O., Toktas H., Demirdal U.S., Subasi V., Kavuncu V., Evcik D. (2014), *Effect of aquatic exercise on ankylosing spondylitis: A randomized controlled trial*, in «Rheumatol. Int.», 34, pp. 1505-1511.
- Eppeland S.G., Diamantopoulos A.P., Soldal D.M., Haugeberg G. (2013), *Short term in-patient rehabilitation in axial spondyloarthritis: The results of 2-week program performed in daily clinical practice*, in «BMC Res Notes», 6, p. 185.
- Eversden L., Maggs F., Nightingale P., Jobanputra P. (2007), *A pragmatic randomised controlled trial of hydrotherapy and land exercises on overall well being and quality of life in rheumatoid arthritis*, in «BMC Musculoskelet. Disord.», 1, pp. 8-23.
- Fernández-de-Las-Peñas C., Alonso-Blanco C., Morales-Cabezas M., Miangolarra-Page J.C. (2005), *Two exercise interventions for the management of patients with ankylosing spondylitis: A randomized controlled trial*, in «Am. J. Phys. Med. Rehabil.», 84, pp. 407-419.

- Gemignani G., Olivieri I., Rujju G., Pasero G. (1991), *Transcutaneous electrical nerve stimulation in ankylosing spondylitis: A double-blind study*, in «Arthr. Rheum.», 34, pp. 788-789.
- Gurcay E., Yuzer S., Eksioğlu E., Bal A., Cakci A. (2008), *Stanger bath therapy for ankylosing spondylitis: Illusion or reality?*, in «Clin. Rheumatol.», 27, pp. 913-917.
- Helliwell P., Abbott C.A., Chamberlain M.A. (1996), *A randomised trial of three different physiotherapy regimes in ankylosing spondylitis*, in «Physiother.», 82, pp. 85-90.
- Hidding A., Van der Linden S., Boers M., Gielen X., De Witte L., Kester A. et al. (1993), *Is group physical therapy superior to individualized therapy in ankylosing spondylitis? A randomized controlled trial*, in «Arthr. Care Res.», 6, pp. 117-125.
- Ince G., Sarpel T., Durgun B., Erdogan S. (2006), *Effects of a multimodal exercise program for people with ankylosing spondylitis*, in «Phys. Ther.», 86, pp. 924-935.
- Karapolat H., Akkoc Y., Sari I., Eyigor S., Akar S., Kirazli Y., Akkoc N. (2008), *Comparison of group-based exercise versus home-based exercise in patients with ankylosing spondylitis: Effects on Bath Ankylosing Spondylitis Indices, quality of life and depression*, in «Clin. Rheumatol.», 27, pp. 695-700.
- Kraag G., Stokes B., Groh J., Helewa A., Goldsmith C. (1990), *The effects of comprehensive home physiotherapy and supervision on patients with ankylosing spondylitis: A randomized controlled trial*, in «J. Rheumatol.», 17, pp. 228-233.
- Lee E.N., Kim Y.H., Chung W.T., Lee M.S. (2008), *Tai Chi for disease activity and flexibility in patients with ankylosing spondylitis: A controlled clinical trial*, in «Evid. Based Complem. Altern. Med.», 5, pp. 457-462.
- Lerner L.A. (1988), *Effectiveness of laser therapy in Bechterew's disease*, in «Ter. Arkh.», 60, pp. 134-136.
- Lim H.J., Moon Y.I., Lee M.S. (2005), *Effects of home-based daily exercise therapy on joint mobility, daily activity, pain, and depression in patients with ankylosing spondylitis*, in «Rheumatol. Int.», 25, pp. 225-229.
- Lubrano E. et al. (2006), *Effects of a combination treatment of an intensive rehabilitation program and etanercept in patients with ankylosing spondylitis: A pilot study*, in «J. Rheumatol.», 33(10), pp. 2029-2034.
- Lubrano E., Spadaro A., Parsons W.J., Atteno M., Ferrara N. (2009), *Rehabilitation in psoriatic arthritis*, in «J. Rheumatol.», 83, pp. 81-82.
- Maddali Bongli S. (2007), *Riabilitazione reumatologica. Approccio multidisciplinare*, Edra, Milano.
- Maddali Bongli S., Del Rosso A. (2010), *How to prescribe physical exercise in rheumatology*, in «Reumat.», 62, pp. 4-11.
- Masiero S., Poli P., Bonaldo L., Pigatto M., Ramonda R., Lubrano E. et al. (2013), *Supervised training and home-based rehabilitation in patients with stabilized ankylosing spondylitis on TNF inhibitor treatment: A controlled clinical trial with a 12-month follow-up*, in «Clin. Rehabil.», 27-28, pp. 562-572.
- McLeod C., Bagust A., Boland A., Dagenais P., Dickson R., Dundar Y. et al. (2007), *Adalimumab, etanercept and infliximab for the treatment of ankylosing spondylitis: A systematic review and economic evaluation*, in «Health Technol. Assess.», 11, pp. 1-158.
- Mihai B., Van der Linden S., De Bie R., Stucki G. (2005), *Experts' beliefs on physiotherapy for patients with ankylosing spondylitis and assessment of their knowledge on published evidence in the field. Results of a questionnaire among international ASAS members*, in «Eur. Medicophys.», 41, pp. 149-153.
- Minor M.A. (2003), *2002 Exercise and physical activity Conference, St. Louis, Missouri: Exercise and arthritis "we know a little bit about a lot of things em leader"*, in «Arthr. Rheum.», 15(49), pp. 1-2.
- Oosterveld F.G., Rasker J.J., Floors M., Landkroon R., Van Rennes B., Zwijnenberg J. et al. (2009), *Infrared sauna in patients with rheumatoid arthritis and ankylosing spondylitis. A pilot study showing good tolerance, short-term improvement of pain and stiffness, and a trend towards long-term beneficial effects*, in «Clin. Rheumatol.», 28, pp. 29-34.

- Ortancil O., Sarikaya S., Sapmaz P., Basaran A., Ozdolap S. (2009), *The effect(s) of a six-week home-based exercise program on the respiratory muscle and functional status in ankylosing spondylitis*, in «J. Clin. Rheumatol.», 15, pp. 68-70.
- Roşu M.O., Ţopa I., Chiriac R., Ancuta C. (2014), *Effects of Pilates, McKenzie and Heckscher training on disease activity, spinal motility and pulmonary function in patients with ankylosing spondylitis: A randomized controlled trial*, in «Rheumatol. Int.», 34, pp. 367-372.
- Spadaro A., De Luca T., Massimiani M.P., Ceccarelli F., Riccieri V., Valesini G. (2008), *Occupational therapy in ankylosing spondylitis: Short-term prospective study in patients treated with anti-TNF- α drugs*, in «Joint Bone Spin.», 75, pp. 29-33.
- Stenström C.H., Arge B., Sundbom A. (1997), *Home exercise and compliance in inflammatory rheumatic diseases: A prospective clinical trial*, in «J. Rheumatol.», 24, pp. 470-476.
- Sweeney S., Taylor G., Calin A. (2002), *The effect of a home based exercise intervention package on outcome in ankylosing spondylitis: A randomized controlled trial*, in «J. Rheumatol.», 29(4), pp. 763-766.
- Van Tubergen A., Landewé R., Van der Heijde D., Hidding A., Wolter N., Asscher M. et al. (2001), *Combined SpA-exercise therapy is effective in patients with ankylosing spondylitis: A randomized controlled trial*, in «Arthr. Rheum.», 45, pp. 430-438.
- Viitanen J.V., Suni J., Kautiainen H., Limatainen M., Takala H. (1992), *Effect of physiotherapy on spinal mobility in ankylosing spondylitis*, in «Scandin. J. Rheum.», 21, pp. 38-41.
- Widberg K., Karimi H., Hafström I. (2009), *Self- and manual mobilization improves spine mobility in men with ankylosing spondylitis: A randomized study*, in «Clin. Rehabil.», 23, pp. 599-608.
- Yurtkuran M., Ay A., Karakoç Y. (2005), *Improvement of the clinical outcome in ankylosing spondylitis by balneotherapy*, in «Joint Bone Spin.», 72, pp. 303-308.